

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN8: Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet	87.6 (81.4-91.9)	928	87.0 (70.3-95.0)	65			●
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	4.5 (3.0-6.6)	1,382	5.2 (2.2-12.2)	111			●
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	20.7 (17.9-23.8)	1,380	29.2 (18.4-43.1)	113			●
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	6.6 (4.7-9.1)	559	11.7 (3.0-35.9)	38			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	8.4 (6.5-10.7)	1,375	12.9 (7.4-21.5)	108			●
QN14: Percentage of students who carried a gun on one or more of the past 30 days	2.6 (1.5-4.4)	1,379	5.8 (2.3-13.6)	109			●
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	1.9 (1.3-2.7)	1,381	2.5 (0.5-12.3)	111			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	5.2 (3.9-6.8)	1,389	10.6 (6.1-17.8)	113			●
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	5.0 (3.4-7.2)	1,363	10.6 (6.1-17.8)	110			●
QN18: Percentage of students who were in a physical fight one or more times during the past 12 months	20.9 (18.0-24.1)	1,366	31.8 (20.5-45.7)	109			●
QN19: Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse	2.3 (1.6-3.3)	1,377	2.4 (0.7-8.2)	109			●
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	8.2 (6.4-10.5)	1,369	13.6 (7.4-23.4)	111			●
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	7.0 (5.0-9.7)	1,384	17.8 (11.5-26.6)	112		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	6.2 (3.9-9.5)	834	16.9 (10.9-25.3)	77		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	9.4 (7.3-12.0)	827	27.6 (17.9-40.0)	77		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	13.0 (9.9-17.0)	1,388	19.8 (13.5-28.1)	111			●
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	6.9 (4.9-9.7)	1,387	19.4 (10.8-32.4)	112		●	
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	25.6 (23.0-28.4)	1,387	45.6 (36.4-55.1)	110		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	10.6 (8.4-13.3)	1,381	29.6 (21.2-39.6)	110		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	9.6 (7.9-11.5)	1,386	33.1 (23.5-44.2)	110		●	
QN29: Percentage of students who actually attempted suicide one or more times during the past 12 months	6.3 (4.6-8.5)	1,191	23.3 (14.9-34.5)	95		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN30: Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse	2.4 (1.7-3.4)	1,177	5.8 (2.9-11.5)	93			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who ever tried cigarette smoking, even one or two puffs	31.5 (27.9-35.4)	1,317	48.1 (36.4-60.1)	106		●	
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	5.6 (3.8-8.0)	1,301	17.1 (11.2-25.3)	106		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	5.5 (3.8-7.7)	1,316	17.2 (10.4-27.1)	101		●	
QNFRDIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	0.5 (0.2-1.4)	1,316	2.8 (1.1-6.8)	101			●
QN34: Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days	0.0 (.-)	77	-	19			
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	17.6 (7.7-35.6)	51	-	10			
QN36: Percentage of students who smoked cigarettes on school property on one or more of the past 30 days	0.9 (0.5-1.5)	1,364	4.5 (1.7-11.1)	105			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN37: Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days	2.7 (1.8-3.9)	1,329	8.8 (4.0-18.2)	106			●
QN38: Among students who reported current cigarette use, the percentage who did not ever try to quit smoking cigarettes during the past 12 months	55.9 (38.2-72.2)	75	-	19			
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	1.6 (1.0-2.4)	1,372	4.5 (2.0-9.7)	113			●
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	5.1 (3.9-6.7)	1,387	14.4 (10.1-20.3)	113		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	7.2 (5.1-9.9)	1,299	18.2 (10.8-29.2)	101		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	58.2 (54.3-62.0)	1,334	81.5 (72.2-88.1)	104		●	
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	16.4 (14.2-18.9)	1,359	31.6 (21.9-43.2)	113		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	24.9 (22.1-27.9)	1,236	59.7 (41.3-75.8)	85		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	11.7 (9.7-14.1)	1,328	28.2 (17.1-42.9)	105		●	
QN45: Percentage of students who had ten or more drinks of alcohol in a row, that is, within a couple of hours, during the past 30 days	2.3 (1.6-3.3)	1,249	5.3 (1.3-19.9)	88			●
QN46: Among students who reported current alcohol use, the percentage who usually got the alcohol they drank from someone who gave it to them during the past 30 days	34.5 (25.8-44.3)	314	35.8 (25.1-48.1)	51			●
QN47: Percentage of students who used marijuana one or more times during their life	37.3 (32.0-42.8)	1,338	64.3 (54.6-73.0)	103		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN48: Percentage of students who tried marijuana for the first time before age 13 years	7.7 (5.6-10.4)	1,367	21.7 (14.4-31.4)	104		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	18.6 (14.7-23.2)	1,373	39.6 (29.4-50.9)	109		●	
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	5.2 (4.0-6.8)	1,384	17.1 (10.1-27.5)	112		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	8.9 (7.2-10.8)	1,368	23.7 (18.2-30.3)	112		●	
QN52: Percentage of students who used heroin one or more times during their life	1.5 (0.9-2.3)	1,385	11.3 (6.4-19.0)	113		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	4.1 (2.6-6.6)	1,382	11.5 (6.5-19.6)	113		●	
QN54: Percentage of students who used ecstasy one or more times during their life	8.4 (6.1-11.5)	1,375	34.0 (23.8-45.9)	110		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	2.1 (1.2-3.6)	1,390	11.8 (8.1-17.0)	111		●	
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	9.2 (6.8-12.5)	1,389	20.0 (12.9-29.5)	111		●	
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	1.2 (0.7-2.0)	1,379	9.4 (4.5-18.6)	112		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	28.6 (25.6-31.9)	1,387	39.8 (31.5-48.7)	106		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	31.4 (26.2-37.1)	1,262	52.1 (33.8-69.9)	87		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	3.5 (2.6-4.8)	1,263	6.7 (3.5-12.5)	88			●
QN61: Percentage of students who had sexual intercourse with four or more people during their life	6.9 (5.3-9.0)	1,261	14.3 (6.5-28.7)	87			●
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	17.5 (15.1-20.2)	1,257	36.7 (24.5-50.9)	87		●	
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	15.0 (9.1-23.7)	218	27.2 (14.8-44.7)	33			●
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	34.7 (30.0-39.8)	213	39.3 (20.8-61.4)	33			●
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	92.9 (88.0-95.9)	211	89.6 (69.9-97.0)	32			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	97.9 (93.0-99.4)	211	96.4 (74.4-99.6)	32			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	90.8 (84.9-94.5)	211	86.0 (61.0-96.0)	32			●
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	98.0 (94.2-99.3)	210	95.1 (81.2-98.9)	32			●
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	17.6 (11.6-25.9)	211	38.1 (20.8-59.0)	32		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) <sup>§</sup>	18.3 (15.7-21.3)	1,277	14.2 (7.5-25.3)	104			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) <sup>§</sup>	13.0 (10.4-16.0)	1,277	21.5 (14.5-30.8)	104		●	
QN66: Percentage of students who described themselves as slightly or very overweight	35.3 (31.5-39.2)	1,369	37.9 (29.1-47.4)	112			●
QN68: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	9.6 (7.9-11.6)	1,363	14.8 (6.8-29.1)	106			●
QN69: Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	4.9 (3.7-6.4)	1,372	9.2 (4.3-18.4)	108			●
QN70: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	4.4 (3.6-5.3)	1,358	9.1 (5.2-15.4)	104			●
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	18.6 (14.7-23.3)	1,350	19.1 (12.8-27.5)	110			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

<sup>§</sup>Based on reference data from the 2000 CDC Growth Charts.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QN72: Percentage of students who did not eat fruit during the past seven days	8.7 (7.1-10.6)	1,366	11.7 (6.5-20.0)	110			●
QN73: Percentage of students who did not eat green salad during the past seven days	37.5 (34.3-40.9)	1,361	35.0 (26.4-44.6)	111			●
QN74: Percentage of students who did not eat potatoes during the past seven days	37.9 (34.9-41.0)	1,354	38.1 (27.2-50.3)	110			●
QN75: Percentage of students who did not eat carrots during the past seven days	45.4 (41.3-49.6)	1,359	48.1 (37.9-58.5)	109			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	20.4 (17.2-24.0)	1,350	24.7 (17.8-33.2)	109			●
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	79.2 (76.0-82.0)	1,320	76.3 (66.2-84.2)	105			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	67.6 (64.2-70.8)	1,348	70.8 (58.9-80.4)	109			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	4.1 (3.0-5.6)	1,348	4.7 (1.7-12.7)	109			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	36.7 (32.2-41.5)	1,348	43.8 (35.1-52.8)	109			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	78.4 (75.9-80.7)	1,348	78.2 (66.7-86.5)	109			●
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	87.1 (85.1-88.9)	1,338	83.9 (73.5-90.7)	107			●
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	7.9 (6.7-9.3)	1,338	5.8 (2.3-13.5)	107			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	43.4 (39.8-47.1)	1,338	40.2 (31.5-49.6)	107			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	76.9 (74.6-79.0)	1,338	74.7 (63.5-83.3)	107			●
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	90.8 (89.0-92.3)	1,320	89.7 (79.6-95.1)	105			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	16.2 (13.3-19.5)	1,342	18.1 (11.1-28.1)	110			●
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	76.2 (72.8-79.4)	1,342	80.1 (71.0-86.9)	110			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	10.1 (8.1-12.4)	1,342	11.0 (4.6-24.1)	110			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	4.9 (3.3-7.3)	1,342	7.3 (3.2-15.6)	110			●
QN78: Percentage of students who did not drink milk or who drank less than three glasses per day of milk during the past seven days	90.5 (87.4-92.9)	1,353	93.3 (84.5-97.3)	109			●
QNMILK0: Percentage of students who drank 0 glasses per day of milk during the past seven days	16.4 (13.4-20.0)	1,353	26.6 (19.4-35.4)	109		●	
QNMILK1: Percentage of students who drank less than one glass per day of milk during the past seven days	64.7 (60.9-68.3)	1,353	74.5 (64.2-82.6)	109		●	
QNMILK2: Percentage of students who drank less than two glasses per day of milk during the past seven days	79.4 (75.6-82.7)	1,353	88.3 (80.2-93.4)	109		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who did not eat breakfast on all of the past seven days	62.7 (58.9-66.3)	1,341	79.2 (69.5-86.5)	109		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	13.8 (11.6-16.3)	1,341	16.3 (9.9-25.6)	109			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	53.9 (49.6-58.1)	1,322	64.8 (51.7-76.1)	109			●
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	14.0 (11.7-16.7)	1,322	14.8 (7.4-27.4)	109			●
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	75.8 (72.1-79.2)	1,322	86.5 (78.8-91.7)	109		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	34.6 (30.8-38.6)	1,340	30.7 (24.4-37.9)	107			●
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	36.3 (33.6-39.2)	1,339	44.6 (34.4-55.4)	108			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	38.8 (27.9-51.0)	1,321	42.4 (30.2-55.6)	107			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	65.7 (51.9-77.4)	1,321	64.2 (46.4-78.7)	107			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN84: Percentage of students who did not play on sports teams during the past 12 months	49.9 (45.9-53.9)	1,334	57.6 (47.8-66.8)	109			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sexual Identity Report

Other	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
Health Risk Behavior							
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	19.1 (14.8-24.3)	1,337	26.0 (20.3-32.6)	105			●
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	18.7 (16.6-20.9)	1,340	22.7 (15.9-31.3)	108			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who have ever been the victim of teasing or name calling during the past 12 months because someone thought they were gay, lesbian, or bisexual	7.3 (5.7-9.3)	1,386	26.4 (16.5-39.3)	110		●	
QN88: Percentage of students who did something to purposely hurt themselves without wanting to die, such as cutting or burning themselves on purpose, one or more times during the past 12 months	13.2 (11.0-15.7)	1,386	34.6 (27.6-42.3)	113		●	
QN89: Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days	7.3 (5.7-9.3)	1,387	14.6 (10.9-19.2)	113		●	
QN90: Percentage of students who used marijuana on school property one or more times during the past 30 days	8.8 (6.0-12.5)	1,377	19.9 (12.5-30.1)	111		●	
QN91: Percentage of students who used any form of cocaine, including powder, crack, or freebase, one or more times during the past 30 days	2.3 (1.5-3.6)	1,381	8.2 (5.3-12.6)	113		●	
QN92: Percentage of students who have taken over-the-counter drugs to get high one or more times during their life	5.8 (4.1-8.0)	1,387	13.3 (7.0-23.7)	112			●
QN93: Percentage of students who have ever had oral sex	27.2 (22.5-32.6)	1,341	44.4 (32.5-56.9)	106		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN94: Percentage of students who have ever had anal sex	9.2 (7.5-11.3)	1,344	18.1 (9.3-32.3)	105			●
QN98: Percentage of students who describe themselves as transgender	6.1 (4.3-8.7)	1,318	9.5 (4.9-17.6)	104			●
QN99: Percentage of students who have not ever been tested or are not sure if they have ever been tested for HIV, the virus that causes AIDS	86.5 (84.4-88.3)	1,333	73.2 (66.3-79.2)	105	●		
QN100: Percentage of students who had been told by a doctor or nurse that they had asthma and who still have asthma	6.5 (4.5-9.3)	1,332	11.3 (7.0-17.9)	107		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.